

A lattice bosonic model as a quantum theory of gravity

Zheng-Cheng Gu^{†,††} and Xiao-Gang Wen[†]

*Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139[†]
Center for Advanced Study, Tsinghua University, Beijing, China, 100084^{††}*

(Dated: February 6, 2008)

A local quantum bosonic model on a lattice is constructed whose low energy excitations are gravitons described by linearized Einstein action. Thus the bosonic model is a quantum theory of gravity, at least at the linear level. We find that the compactification and the discretization of metric tensor are crucial in obtaining a quantum theory of gravity.

Seven wonders of our universe: Our world has many mysteries and wonders. At the most fundamental level, there are may be seven deep mysteries and/or wonders in our universe: (1) Identical particles.¹⁻³ (2) Gauge interactions.¹⁻³ (3) Fermi statistics.^{4,5} (4) Tiny masses of fermions ($\sim 10^{-20}$ of the Planck mass).^{6,7,12} (5) Chiral fermions.^{8,9} (6) Lorentz invariance.¹⁰ (7) Gravity.¹¹ Here we would like to question where those wonderful and mysterious properties come from. We wish to have a single unified understanding of all of the above mysteries. Or more precisely, we wish that we can start from a single model to obtain all of the above wonderful properties.

Recent progresses showed that, starting from a single origin – a local bosonic model (which is also called spin model), the first four properties in the above list emerge at low energies.¹²⁻¹⁵ Thus we may say that the local bosonic model provides a unified origin for identical particles, gauge interactions, Fermi statistics and near masslessness of the fermions. Gauge interaction and Fermi statistics are unified under this point of view of emergence. However, three more mysteries remain to be understood. In this paper, we will show that it is possible to construct a local bosonic model from which gravitons emerge. On one hand, such a model can be viewed as a theory of quantum gravity. It solves a long standing problem of putting quantum mechanics and gravity together. On the other hand, the model provides a design of a condensed matter system which has an emergent quantum gravity at low energies. This allows us to study some quantum effects of gravity (a simulated one) in a laboratory.

The belief that all the wonderful phenomena of our universe (such as gauge interaction, Fermi statistics, and gravitons) emerge from a lowly local bosonic model is called locality principle.¹² Using local bosonic model as a underlying structure to understand the deep mysteries of our universe represent a departure from the traditional approach of gaining a better understanding by dividing things in to smaller parts. In the traditional approach, we assume that the division cannot continue forever and view everything as made of some simple indivisible building blocks – the elementary particles. However, the traditional approach may represent a wrong direction. For example, phonons behave just like any other elementary particles at low energies. But if we look at phonons closely, we do not see smaller parts that form a

phonon. We see the atoms that form the entire lattice. The phonons are not formed by those atoms, the phonons are simply collective motions of those atoms. This makes us to wonder that photons, electrons, gravitons, *etc*, may also be emergent phenomena just like phonons. They may not be the building blocks of everything. They may be collective motions of a deeper underlying structure.

This paper uses a particular emergence approach: we try to obtain everything from a local bosonic model. The detail form of the bosonic model is not important. The important issue is how the bosons (or the spins) are organized in the ground state. It is shown that if bosons organize into a string-net condensed state, then photons, electrons and quarks can emerge naturally as collective motions of the bosons.¹²⁻¹⁵ In this paper, we will find an organization of bosons such that the collective motions of bosons lead to gravitons.

Other emergence approaches were developed in superstring theory¹⁶ in last 10 years, as demonstrated by the duality relations among various superstring models and matrix models.¹⁷ The anti-de Sitter-space/conformal-field-theory duality even shows how space-time and gravity emerge from a gauge theory.¹⁸

The rule of game: There are many different approaches to quantum gravity²⁰⁻²² based on different principles. Some approaches, such as loop quantum gravity,²³ stress the gauge structure from the diffeomorphism of the space-time. Other approaches, such as superstring theory,^{16,17} stress the renormalizability of the theory. In this paper, we follow a different rule of game by stressing finiteness and locality.

Our rule of game is encoded in the following working definition of quantum gravity. Quantum gravity is

- (a) a quantum theory.
- (b) Its Hilbert space has a finite dimension.
- (c) Its Hamiltonian is a sum of local operators.
- (d) The gapless helicity ± 2 excitations are the only low energy excitations.
- (e) The helicity ± 2 excitations have a linear dispersion.
- (f) The gravitons³⁵ can interact in the way consistent with experimental observations.

The condition (b) implies that the quantum gravity considered here has a finite cut-off. So the renormalizability is not an issue here. The condition (c) is a locality condition. It implies two additional things: (1) the total Hilbert space is a direct product of local Hilbert spaces

$\mathcal{H}_{tot} = \otimes_n \mathcal{H}_n$, (2) the local operators are defined as operators that act within each local Hilbert space $\mathcal{H}_n$ or finite products of local operators. The conditions (a – c) actually define a local bosonic model. Certainly, any quantum spin models satisfy (a – c). It is the conditions (d–f) that make the theory to look like gravity.

The condition (d) is very important. It is very easy to construct a quantum model that contains helicity ± 2 gapless excitations, such as the theory described by the following Lagrangian $\mathcal{L} = \frac{1}{2} \partial_0 h^{ij} \partial_0 h^{ij} - \frac{1}{2} \partial_k h^{ij} \partial_k h^{ij}$. Such a theory is not a theory of gravity since it also contain helicity 0, ± 1 gapless excitations.

Superstring theory^{16,17} satisfies the conditions (a), (e) and (f), but in general not (d) due to the presence of dilatons (massless scalar particles). The superstring theory (or more precisely, the superstring field theory) also does not satisfy the condition (b) since the cut-off is not explicitly implemented. The spin network²⁴ or the quantum computing²⁵ approach to quantum gravity satisfies the condition (a,b) or (a–c). But the properties (d–f) are remain to be shown. The induced gravity from superfluid ³He discussed in Ref. 26 does not satisfy the condition (d) due to the presence of gapless density mode. In Ref. 27, it is proposed that gravitons may emerge as edge excitations of a quantum Hall state in 4 spatial dimensions. Again the condition (d) is not satisfied due to the presence of infinite massless helicity $\pm 1, \pm 2, \pm 3, \dots$ modes.

In Ref. 28, a very interesting bosonic model is constructed which contains gapless helicity 0 and ± 2 excitations with quadratic dispersions. The model satisfies (a–c), but not (d,e). In this paper, we will fix the two problems and construct a local bosonic model that satisfies the conditions (a – e) and possibly (f). To quadratic order, the low energy effective theory of our model is the linearized Einstein gravity. The key step in our approach is to *discretize and compactify* the metric tensor.

Review of emergence of $U(1)$ gauge theory: Our model for emergent quantum gravity is closely related to the rotor model that produces emergent $U(1)$ gauge theory.^{19,30} So we will first discuss the emergence of $U(1)$ gauge theory to explain the key steps in our argument in a simpler setting.

To describe the rotor model, We introduce an angular variable $a_{ij} \sim a_{ij} + 2\pi$ and the corresponding angular momentum $\mathcal{E}_{ij}$ for each link of a cubic lattice. Here i labels the sites of the cubic lattice and a_{ij} and $\mathcal{E}_{ij}$ satisfy $a_{ij} = -a_{ji}$ and $\mathcal{E}_{ij} = -\mathcal{E}_{ji}$. The phase space Lagrangian for physical degrees of freedom a_{ij} and $\mathcal{E}_{ij}$ is given by¹⁹

$$L = \sum_{\langle ij \rangle} \mathcal{E}_{ij} \dot{a}_{ij} - \frac{J}{2} \sum_{\langle ij \rangle} \mathcal{E}_{ij}^2 + g \sum_{\langle ijkl \rangle} \cos B_{ijkl} - \frac{U}{2} \sum_i Q_i^2$$

$$B_{ijkl} = a_{ij} + a_{jk} + a_{kl} + a_{li}, \quad Q_i = \sum_{j \text{ next to } i} \mathcal{E}_{ij} \quad (1)$$

where $\sum_i$ sums over all sites, $\sum_{\langle ij \rangle}$ over all links, and $\sum_{\langle ijkl \rangle}$ over all square faces of the cubic lattice. We note that after quantization, $\mathcal{E}_{ij}$ are quantized as integers.

To obtain the low energy dynamics of the above rotor model, let us assume that the fluctuation of $\mathcal{E}_{ij}$ are large and treat $\mathcal{E}_{ij}$ as a continuous quantity. We also assume that the fluctuations of a_{ij} are small and expand (1) to the quadratic order of a_{ij} . Then we take the continuum limit by introducing the continuous fields $(\mathcal{E}^i, a^i)$ and identifying $\mathcal{E}_{ij} = \int_i^j dx^i \mathcal{E}^i$ and $a_{ij} = \int_i^j dx^i a^i$. Here we have assumed that the lattice constant $a = 1$. The resulting continuum effective theory is given by

$$\mathcal{L} = -\mathcal{E}^i \partial_0 a_i - \frac{1}{2} J (\mathcal{E}^i)^2 - \frac{1}{2} g (\mathcal{B}^i)^2 - \frac{1}{2} U (\partial_i \mathcal{E}^i)^2, \quad (2)$$

where $\mathcal{B}^i = \epsilon^{ijk} \partial_j a_k$. We find that the rotor model has three low lying modes. Two of them are helicity ± 1 modes with a linear dispersion $\omega_{\mathbf{k}} \sim \sqrt{gJ} |\mathbf{k}|$ and the third mode is the helicity 0 mode with zero frequency $\omega_{\mathbf{k}} = 0$. We know that a $U(1)$ gauge theory only have two helicity ± 1 modes at low energies. Thus the key to understand the emergence $U(1)$ gauge theory is to understand how the helicity 0 mode obtain an energy gap.

To understand why helicity 0 mode is gapped, let us consider the quantum fluctuations of $\mathcal{E}^i$ and a_i . We note that the longitudinal mode and the transverse modes separate. Introduce $\mathcal{E}^i = \mathcal{E}_{||} + \mathcal{E}_{\perp}$ and $a_i = a_{||} + a_{\perp}$, we find that the dynamics of the transverse mode is described by $\mathcal{L}_{\perp} = a_{\perp} \partial_0 \mathcal{E}_{\perp} - \frac{J}{2} \mathcal{E}_{\perp}^2 - \frac{g}{2} \partial_i a_{\perp} \partial_i a_{\perp}$. At the lattice scale $\delta x \sim 1$, the quantum fluctuations of $\mathcal{E}_{\perp}$ and $a_{\perp}$ are given by $\delta \mathcal{E}_{\perp} \sim \sqrt{\frac{J}{g}}$ and $\delta a_{\perp} \sim \sqrt{\frac{g}{J}}$. We see that the assumptions that we used to derive the continuum limit are valid when $J \ll g$. In this limit we can trust the result from the continuum effective theory and conclude that the transverse modes (or the helicity ± 1 modes) have a linear gapless dispersion.

The longitudinal mode is described by $(f(\mathbf{x}), \pi(\mathbf{x}))$ with $a_i = \partial_i f$ and $\pi = \partial_i \mathcal{E}^i$. Its dynamics is determined by $\mathcal{L}_{||} = \pi \partial_0 f - \frac{J}{2} \pi (-\partial^{-2}) \pi - \frac{1}{2} U \pi^2$. At the lattice scale, the quantum fluctuations of π and f are given by $\delta \pi = 0$ and $\delta f = \infty$. We see that a positive U and J will make the fluctuations of f much bigger than the compactification size 2π and the fluctuations of π much less than the discreteness of $\mathcal{E}^i$ which is 1. In this limit, the result from the classical equation of motion cannot be trusted.

In fact the weak quantum fluctuations in the discrete variable π and the strong quantum fluctuations in the compact variable f indicate that the corresponding mode is gapped after the quantization. Since π has weak fluctuations which is less than the discreteness of π , the ground state is basically given by $\pi = 0$. A low lying excitation is then given by $\pi = 0$ everywhere except in a unit cell where $\pi = 1$. Such an excitation have an energy of order U . The gapping of helicity 0 mode is confirmed by more careful calculations.¹⁹ From those calculations, we find that the weak fluctuations of π lead to a constraint $\pi = \partial_i \mathcal{E}^i = 0$ and the strong fluctuations of f lead to a gauge transformation $a_i \rightarrow a_i + \partial_i f$. The Lagrangian (2) equipped with the above constraint and the gauge transformation becomes the Lagrangian of a $U(1)$ gauge

theory. We will use this kind of argument to argue the emergence of gravitons and the linearized Einstein action.

The emergence of quantum gravity: First, let us describe a model that will have emergent gravitons at low energies. The model has six variables $\theta_{xx}(\mathbf{i})$, $\theta_{yy}(\mathbf{i})$, $\theta_{zz}(\mathbf{i})$, $L^{xx}(\mathbf{i})$, $L^{yy}(\mathbf{i})$, and $L^{zz}(\mathbf{i})$ on each vertex of a cubic lattice. The model also has two variables on each square face of the cubic lattice. For example, on the square centered at $\mathbf{i} + \frac{\mathbf{x}}{2} + \frac{\mathbf{y}}{2}$, the two variables are $\theta_{xy}(\mathbf{i} + \frac{\mathbf{x}}{2} + \frac{\mathbf{y}}{2}) = \theta_{yx}(\mathbf{i} + \frac{\mathbf{x}}{2} + \frac{\mathbf{y}}{2})$ and $L^{xy}(\mathbf{i} + \frac{\mathbf{x}}{2} + \frac{\mathbf{y}}{2}) = L^{yx}(\mathbf{i} + \frac{\mathbf{x}}{2} + \frac{\mathbf{y}}{2})$. The bosonic model is described by the following phase space Lagrangian

$$L = \sum_{\mathbf{i}, ab=xy,yz,zx} L^{ab}(\mathbf{i} + \frac{\mathbf{a}}{2} + \frac{\mathbf{b}}{2}) \partial_0 \theta_{ab}(\mathbf{i} + \frac{\mathbf{a}}{2} + \frac{\mathbf{b}}{2}) + \sum_{\mathbf{i}, a=x,y,z} L^{aa}(\mathbf{i}) \partial_0 \theta_{aa}(\mathbf{i}) - H_U - H_J - H_g \quad (3)$$

where

$$\begin{aligned} H_U &= n_G U_1 \sum_{\mathbf{i}} \sum_{a=x,y,z} \{1 - \cos[2\pi Q(\mathbf{i}, \mathbf{i} + \mathbf{a})/n_G]\} \\ &\quad + n_G U_2 \sum_{\mathbf{i}} \{1 - \cos[\eta(\mathbf{i})]\} \\ H_J &= n_G J \sum_{\mathbf{i}, a=x,y,z} \{1 - \cos[2\pi L^{aa}(\mathbf{i})/n_G]\} \\ &\quad + 2n_G J \sum_{\mathbf{i}} \sum_{ab=xy,yz,zx} \{1 - \cos[2\pi L^{ab}(\mathbf{i} + \frac{\mathbf{a}}{2} + \frac{\mathbf{b}}{2})/n_G]\} \\ &\quad - \frac{1}{2} n_G J \sum_{\mathbf{i}} \{1 - \cos[2\pi \sum_{a=x,y,z} L^{aa}(\mathbf{i})/n_G]\} \\ H_g &= \frac{n_G g}{4} \sum_{\mathbf{i}, a=x,y,z} \{1 - \cos[\rho_a^a(\mathbf{i})]\} \\ &\quad + \frac{n_G g}{4} \sum_{\mathbf{i}, ab=xy,yz,zx} \sin[\rho_b^a(\mathbf{i})] \sin[\rho_a^b(\mathbf{i})] \end{aligned} \quad (4)$$

Here $\rho_j^i(\mathbf{i})$, $\eta(\mathbf{i})$ and $Q(\mathbf{i}, \mathbf{i} + \mathbf{x})$ are defined as $\rho_x^x(\mathbf{i}) = \theta_{zx}(\mathbf{i} + \mathbf{y} + \frac{\mathbf{z}}{2} + \frac{\mathbf{x}}{2}) + \theta_{zx}(\mathbf{i} + \frac{\mathbf{z}}{2} + \frac{\mathbf{x}}{2}) - \theta_{xy}(\mathbf{i} + \mathbf{z} + \frac{\mathbf{x}}{2} + \frac{\mathbf{y}}{2}) - \theta_{xy}(\mathbf{i} + \frac{\mathbf{x}}{2} + \frac{\mathbf{y}}{2})$, $\rho_y^x(\mathbf{i}) = -\theta_{yz}(\mathbf{i} + \frac{\mathbf{z}}{2} + \frac{\mathbf{y}}{2}) - \theta_{yz}(\mathbf{i} + \frac{\mathbf{z}}{2} - \frac{\mathbf{y}}{2}) + 2\theta_{yy}(\mathbf{i} + \mathbf{z}) + 2\theta_{yy}(\mathbf{i})$, $\rho_z^x(\mathbf{i}) = -2\theta_{zz}(\mathbf{i} + \mathbf{y}) - 2\theta_{zz}(\mathbf{i}) + \theta_{yz}(\mathbf{i} + \frac{\mathbf{y}}{2} + \frac{\mathbf{z}}{2}) + \theta_{yz}(\mathbf{i} + \frac{\mathbf{y}}{2} - \frac{\mathbf{z}}{2})$, $\eta(\mathbf{i}) = \sum_{a=x,y,z} \sum_{b=x,y,z} [\theta_{bb}(\mathbf{i} + \mathbf{a}) + \theta_{bb}(\mathbf{i} - \mathbf{a}) + 2\theta_{bb}(\mathbf{i})] - \sum_{a=x,y,z} [\theta_{aa}(\mathbf{i} + \mathbf{a}) + \theta_{aa}(\mathbf{i} - \mathbf{a}) + 2\theta_{aa}(\mathbf{i})] - \sum_{ab=xy,yz,zx} [\theta_{ab}(\mathbf{i} + \frac{\mathbf{a}}{2} + \frac{\mathbf{b}}{2}) + \theta_{ab}(\mathbf{i} - \frac{\mathbf{a}}{2} + \frac{\mathbf{b}}{2}) + \theta_{ab}(\mathbf{i} + \frac{\mathbf{a}}{2} - \frac{\mathbf{b}}{2}) + \theta_{ab}(\mathbf{i} - \frac{\mathbf{a}}{2} - \frac{\mathbf{b}}{2})]$, and $Q(\mathbf{i}, \mathbf{i} + \mathbf{x}) = L^{xx}(\mathbf{i} + \mathbf{x}) + L^{xx}(\mathbf{i}) + L^{yx}(\mathbf{i} + \frac{\mathbf{x}}{2} + \frac{\mathbf{y}}{2}) + L^{yx}(\mathbf{i} + \frac{\mathbf{x}}{2} - \frac{\mathbf{y}}{2}) + L^{zx}(\mathbf{i} + \frac{\mathbf{x}}{2} + \frac{\mathbf{z}}{2}) + L^{zx}(\mathbf{i} + \frac{\mathbf{x}}{2} - \frac{\mathbf{z}}{2})$. Other components are obtained by cycling xyz to yzx and zxy .

Note that both θ_{ab} and its canonical conjugate L^{ab} are compactified: $\theta_{ab} \sim \theta_{ab} + 2\pi$, $L^{ab} \sim L^{ab} + n_G$. Hence after quantization they are both discretized and n_G is an integer. Due to the compactification, only $W_L^{ab} = e^{2\pi i L^{ab}/n_G}$, $W_\theta^{ab} = e^{i\theta_{ab}}$ and their products are physical operators. For a fix ab and $\mathbf{i}$, $W_L^{ab}(\mathbf{i})$ and $W_\theta^{ab}(\mathbf{i})$ satisfy the algebra

$$W_L^{ab}(\mathbf{i}) W_\theta^{ab}(\mathbf{i}) = e^{2\pi i/n_G} W_\theta^{ab}(\mathbf{i}) W_L^{ab}(\mathbf{i}) \quad (5)$$

Such an algebra has only one n_G dimensional representation. This n_G dimensional representation becomes our local Hilbert space $\mathcal{H}_{i,ab}$. The total Hilbert space of our model (3) is given by $\mathcal{H} = \otimes_{i,ab} \mathcal{H}_{i,ab}$ after quantization. In other words, there are n_G^3 states on each vertex and n_G states on each square face of the cubic lattice. Note that the Hamiltonian

$$H = H_U + H_J + H_g \quad (6)$$

is a function of the physical operators W_L^{ab} , W_θ^{ab} and their hermitian conjugates. So the quantum model defined through the Hamiltonian (6) and the algebra (5) is a bosonic model whose local Hilbert spaces have finite dimensions.

Next, we would like to understand low energy excitations of the quantum bosonic model (6) in large n_G limit. We first assume that the fluctuations of $\phi^{ij} \equiv 2\pi L^{ij}/n_G$ and θ_{ij} are much bigger than $1/n_G$ (the discreteness of ϕ^{ij} and θ_{ij}) so that we can treat ϕ^{ij} and θ_{ij} as continuous variables. We also assume that the fluctuations of ϕ^{ij} and θ_{ij} are much smaller than 1 so that we can treat ϕ^{ij} and θ_{ij} as small variables. Under those assumptions, we can use semiclassical approach to understand the low energy dynamics of the quantum bosonic model (6).

Expanding the Lagrangian (3) to quadratic order in ϕ^{ij} and θ_{ij} , we can find the dispersions of collective modes of the bosonic model. There are total of six collective modes. We find four of them have zero frequency for all $\mathbf{k}$, and two modes have a linear dispersion relation near $\mathbf{k} = (\pi, \pi, \pi)$. Near $\mathbf{k} = (\pi, \pi, \pi)$, the dynamics of the six modes are described by the following continuum field theory:

$$\begin{aligned} \mathcal{L} &= n_G \left\{ \phi^{ij} \dot{\theta}_{ij} - \frac{J}{2} [(\phi^{ij})^2 - \frac{(\phi^{ii})^2}{2}] - \frac{g}{2} \theta_{ij} R^{ij} \right. \\ &\quad \left. - \frac{U_1}{2} (\partial_i \phi^{ij})^2 - \frac{U_2}{2} (R^{ii})^2 \right\} \end{aligned} \quad (7)$$

where $R^{ij} = \epsilon^{imk} \epsilon^{jln} \partial_m \partial_l \theta_{nk}$ and we define the continuum field $\theta^{ab}(\mathbf{x})$ as $-(-1)^i \frac{1}{2} \theta^{ab}(\mathbf{i} + \frac{\mathbf{a}}{2} + \frac{\mathbf{b}}{2})$ for $a \neq b$ and as $(-1)^i \theta^{ab}(\mathbf{i})$ for $a = b$. The helicity ± 2 modes have a linear dispersion relation $\omega_{\mathbf{k}} \sim \sqrt{gJ} |\mathbf{k}|$. We find that for large n_G , the quantum fluctuations of the helicity ± 2 modes is of order $\delta\phi^{ij}, \delta\theta_{ij} \sim \sqrt{1/n_G}$ (assuming $U_{1,2}$, J and g are of the same order). So the fluctuations of ϕ^{ij} and θ_{ij} satisfy $1/n_G \ll \delta\phi^{ij}, \delta\theta_{ij} \ll 1$ and the semiclassical approximation is valid for the helicity ± 2 modes. In this case, the result $\omega_{\mathbf{k}} \sim \sqrt{gJ} |\mathbf{k}|$ can be trusted.

The helicity ± 1 modes and one of the helicity 0 mode are described by $\theta_{ij} = \partial_i \theta_j + \partial_j \theta_i$ and $\phi^i = \partial_j \phi^{ji}$. Their frequency $\omega_{\mathbf{k}} = 0$. For such modes, the Hamiltonian only contains ϕ^i . Thus the quantum fluctuations satisfies $\delta\phi^i \ll 1/n_G$ and $\delta\theta_i \gg 1$. So the semiclassical approximation is not valid and the result $\omega_{\mathbf{k}} = 0$ cannot be trusted. Using the similar argument used in emergence of $U(1)$ gauge bosons, we conclude that those modes are gapped. The strong fluctuations $\delta\theta_{ij} = \partial_i \theta_j + \partial_j \theta_i \gg 1$

and the weak fluctuations $\phi^i \ll 1/n_G$ lead to gauge transformations and the constraints

$$\theta_{ij} \rightarrow \theta_{ij} + \partial_i \theta_j + \partial_j \theta_i, \quad \partial_j \phi^{ji} = 0 \quad (8)$$

The second helicity 0 mode is described by $\phi^{ij} = (\delta_{ij}\partial^2 - \partial_i\partial_j)\phi$ and $\theta = (\delta_{ij}\partial^2 - \partial_i\partial_j)\theta_{ij}$. Its frequency is again $\omega_k = 0$. The Hamiltonian for such a mode contains only θ . So the quantum fluctuations satisfies $\delta\phi \gg 1$ and $\delta\theta \ll 1/n_G$. The second helicity 0 mode is also gapped. The strong fluctuations $\delta\phi^{ij} = (\delta_{ij}\partial^2 - \partial_i\partial_j)\phi \gg 1$ and the weak fluctuations $\theta = (\delta_{ij}\partial^2 - \partial_i\partial_j)\theta_{ij} \ll 1/n_G$ lead to a gauge transformation and a constraint

$$\phi^{ij} \rightarrow \phi^{ij} + (\delta_{ij}\partial^2 - \partial_i\partial_j)\phi, \quad (\delta_{ij}\partial^2 - \partial_i\partial_j)\theta_{ij} = 0 \quad (9)$$

The Lagrangian (7) equipped with the gauge transformations and the constraints (8,9) is nothing but the linearized Einstein Lagrangian of gravity, where $\theta_{ij} \sim g_{ij} - \delta_{ij}$ represents the fluctuations of the metric tensor g_{ij} around the flat space. So the linearized Einstein gravity emerge from the quantum model (6) in the large n_G limit. The local bosonic model (6) can be viewed as a quantum theory of gravity.

We have seen that the gapping of the helicity 0 mode in the rotor model (1) leads to an emergence of $U(1)$ gauge structure at low energies. The emergence of a gauge structure also represents a new kind of order – quantum order^{12,31} – in the ground state. In Ref. 32, it was shown that the emergent $U(1)$ gauge invariance, and hence the quantum order, is robust against any local perturbations of the rotor model. Thus the gaplessness of the emergent photon is protected by the quantum order.³³ Similarly, the gapping of the two helicity 0 modes and the

helicity ± 1 modes in the bosonic model (6) leads to an emergent gauge invariance of the linearized coordinate transformation. This indicates that the ground state of the bosonic model contains a new kind of quantum order that is different from those associated with emergent ordinary gauge invariances of internal degrees of freedom. We expect such an emergent linearized diffeomorphism invariance to be robust against any local perturbation of the bosonic model. Thus the gaplessness of the emergent gravitons is protected by the quantum order.

The emergent gravitons in the model (3) naturally interact but the interaction may be different from that described by the higher order non-linear terms in Einstein gravity. However, those higher order terms are irrelevant at low energies. Thus it may be possible to generate those higher order terms by fine tuning the lattice model (3), such as modifying the Hamiltonian (H_J and H_g), the constraints (H_U), as well as the Berry's phase term in (3). So it may be possible that local bosonic models can generate proper non-linear terms to satisfy (f).²⁹

Our result appears to contradict with the Weinberg-Witten theorem³⁴ which states that in all theories with a Lorentz-covariant energy-momentum tensor, composite as well as elementary massless particles with helicity $h > 1$ are forbidden. However, the energy-momentum tensor in our model is not invariant under the linearized diffeomorphism (although the action is invariant). This may be the reason why emergent gravitons are possible in our model.

We would like to thank J. Polchinski and E. Witten for their very helpful comments. This research is supported by NSF grant No. DMR-0433632 and ARO grant No. W911NF-05-1-0474.

¹ H. Weyl, *Space, Time, Matter* (Dover, 1952).

² W. Pauli, *Rev. Mod. Phys.* **13**, 203 (1941).

³ C. N. Yang and R. L. Mills, *Phys. Rev.* **96**, 191 (1954).

⁴ E. Fermi, *Z. Phys.* **36**, 902 (1926).

⁵ P. A. M. Dirac, *Proc. Roy. Soc.* **A112**, 661 (1926).

⁶ D. J. Gross and F. Wilczek, *Phys. Rev. Lett.* **30**, 1343 (1973).

⁷ H. D. Politzer, *Phys. Rev. Lett.* **30**, 1346 (1973).

⁸ T. Lee and C. Yang, *Phys. Rev.* **104**, 254 (1956).

⁹ C. Wu et al., *Phys. Rev.* **105**, 1413 (1957).

¹⁰ A. Einstein, *Annalen der Physik* **17**, 891 (1905).

¹¹ A. Einstein, *Annalen der Physik* **49**, 769 (1916).

¹² X.-G. Wen, *Phys. Rev. D* **68**, 065003 (2003).

¹³ M. A. Levin and X.-G. Wen, *Rev. Mod. Phys.* **77**, 871 (2005).

¹⁴ M. Levin and X.-G. Wen, hep-th/0507118.

¹⁵ X.-G. Wen, cond-mat/0508020.

¹⁶ M. B. Green, J. H. Schwarz, and E. Witten, *Superstring Theory* (Cambridge University Press, 1988).

¹⁷ J. Polchinski, *String Theory* (Cambridge University Press, 1998).

¹⁸ O. Aharony *etal*, *Phys. Rept.* **323**, 183 (2000).

¹⁹ X.-G. Wen, *Quantum Field Theory of Many-Body Systems*

– *From the Origin of Sound to an Origin of Light and Electrons* (Oxford Univ. Press, Oxford, 2004).

²⁰ T. Thiemann, gr-qc/0210094.

²¹ S. Carlip, *Rept. Prog. Phys.* **64**, 885 (2001).

²² P. Peldan, *Class. Quant. Grav.* **11**, 1087 (1994).

²³ L. Smolin, hep-th/0408048.

²⁴ C. Rovelli and L. Smolin, *Phys. Rev.* **D52**, 5743 (1995).

²⁵ L. Smolin, hep-th/0507235.

²⁶ G. Volovik, *J. Low. Temp. Phys.* **113**, 667 (1998).

²⁷ S.-C. Zhang and J. Hu, *Science* **294**, 823 (2001).

²⁸ C. Xu, cond-mat/0602443.

²⁹ Z.-C. Gu and X.-G. Wen, to appear.

³⁰ O. I. Motrunich and T. Senthil, *Phys. Rev. Lett.* **89**, 277004 (2002).

³¹ X.-G. Wen, *Phys. Rev. B* **65**, 165113 (2002).

³² M. B. Hastings and X.-G. Wen, *Phys. Rev B* **72**, 045141 (2005).

³³ X.-G. Wen, *Phys. Rev. Lett.* **88**, 11602 (2002).

³⁴ S. Weinberg and E. Witten, *Phys. Lett. B* **96**, 59 (1980).

³⁵ In this paper, we define gravitons as linearly dispersing gapless helicity ± 2 excitations.